



[Home](#) | [About](#) | [Table of Contents](#) | [Advanced Search](#) | [Copyright](#) | [Feedback](#) | [Privacy](#)

You are here: [Chapter: 1 Units and fundamental constants](#)

Section: 1.2 Fundamental physical constants

SubSection: 1.2.3 Atomic constants

[« Previous Subsection](#)

[Next Subsection »](#)

Unless otherwise stated this page contains Version 1.0 content ([Read more about versions](#))

1.2.3 Atomic constants

The values of e , h , N_A etc., given in this section are based on the values of Cohen and Taylor (1987).

The best values of constants such as the mass and charge of the electron are not found by measuring each separately (e.g. as in Millikan's oil-drop experiment). The values recommended in the above paper were obtained by a weighted 'least-squares' statistical treatment of selected experimental data. The values depend on comparatively few (22) experimental results and the number of 'unknowns' was reduced by treating certain combinations of the constants as being exact (or auxiliary constants) for the purposes of the evaluation. This group included quantities such as the Rydberg constant $\mu_0 c^3 m_e e^4 / 8 h^3$, the ratio μ'_p / μ_B between the proton magnetic moment (in water) μ'_p and the Bohr magneton $\mu_B (= eh/4\pi m_e)$, the Josephson effect value of $2e/h$ in terms of the maintained representation of the volt, and set numerical functional relationships between e , h , and m_e . The unknowns included the spectroscopic fine-structure constant $\alpha ((\mu_0 c^2 / 4\pi) e^2 / h)$, the ratio K_Ω of the maintained representation of the ohm to the SI ohm, the ratio K_V of the BIPM maintained representation of the volt to the SI volt, d_{220} the (220) lattice spacing of a perfect crystal of pure silicon at 22.5 °C in vacuum and μ_μ / μ_p the ratio of the magnetic moment of the muon to that of the proton. Some of the experimental data can only be interpreted in terms of α , etc. by invoking very sophisticated quantum electrodynamic calculations and so, as time passes, improved values are obtained for α as higher order terms are taken into account.

After the 1973 evaluation by Cohen and Taylor it became apparent that the 1973 recommended value of K_V was in error by about eight parts per million. In addition there were further accurate measurements such as the gravitational constant, the gas constant, the Avogadro constant, and the Rydberg constant. The advent of ion traps led to increased accuracy of measurement of the electron magnetic moment anomaly $g_c - 2$, and of the ratio of the proton to electron mass. Better direct realizations were made of the ampere, watt, and ohm, while the quantized Hall resistance in MOSFET semi-conductors at low temperatures provided information concerning K_Ω and α .

As a result of the method of evaluation there are correlations between the output values, so that the full variance and co-variance matrix should be used for computing values that are not given in the table. Although the accuracy is normally well ahead of user requirements, it is always important to specify which evaluation one has used and also to ensure that they come from a consistent set of constants. (Results from different evaluations

may differ and consequently should not be mixed, otherwise the constants will not form a consistent set.) For the most part, more accurate measurements since 1987 have confirmed the correctness of the 1987 set within their assigned uncertainties. A more accurate evaluation will be made when sufficient experimental data are available.

Values accepted formerly. Other conventional values of e , h , etc. may be encountered in books, and some of these earlier values are given below. The first is essentially Millikan's oil-drop value of e , 4.77×10^{-10} esu, and the second the 'X-ray grating' value 4.802. The numbers in brackets are standard errors. It will be seen that the accuracies claimed have sometimes appeared over-optimistic in retrospect, and that besides the supposed random errors of observation some unsuspected systematic error has been present. (In Millikan's value of e , accepted from 1917 to 1935, it was an inaccuracy in the assumed viscosity of air.) Consequently, it may be thought prudent to regard the standard errors given here with caution. Although the accuracy has improved over the years there is no sustained evidence for any systematic time dependence of these constants, and increasing reliance is placed on them for metrological purposes.

<i>Author of discussion</i>	$e \times 10^{19}/C$	$h \times 10^{34}/(J s)$	<i>Kaye and Laby</i>
Birge, 1929.	1.591 1 (0.002 4)	6.547 (0.012)	7th ed.
Birge, 1942.	1.602 03 (0.000 50)	6.624 2 (0.003 5)	10th ed.
Dumond and Cohen, 1963.	1.602 10 (0.000 02)	6.625 6 (0.000 16)	13th ed.
Taylor et al., 1969.	1.602 192 (0.000 007)	6.626 20 (0.000 05)	14th ed.
Cohen and Taylor, 1973.	1.602 189 2 (0.000 004 6)	6.626 176 (0.000 036)	15th ed. (1986)
Cohen and Taylor, 1987.	1.602 177 3 (0.000 000 49)	6.626 075 5 (0.000 004 0)	15th ed. (1991 reprint)

Table of fundamental constants

These are the recommended values of Mohr and Taylor (2004). These are known as the 2002 best values and are based on the published data to about December 2002; they may be superseded by a more accurate set if sufficient new data are available. They will be available on the NIST Web Site at <http://physics.nist.gov/cuu/Constants/>

			<i>SI unit</i>	<i>Standard error</i> (parts in 10^6)
Principal constants				
Speed of light in vacuum	c	2.997 924 580	$\times 10^8 \text{ m s}^{-1}$	exact
Planck constant	h	6.626 0693	$\times 10^{-34} \text{ J s}$	0.17

Planck constant ($h/2\pi$)	\hbar	1.054 571 68	$\times 10^{-34}$ J s	0.17
Elementary charge	e	1.602 176 53	$\times 10^{-19}$ C	0.085
Mass of electron	m_e	9.109 3826	$\times 10^{-31}$ kg	0.17
Mass of electron in atomic mass units		5.485 799 0945	$\times 10^{-4}$ u	0.00044
Avogadro constant	N_A, L	6.022 14145	$\times 10^{23}$ mol ⁻¹	0.17
Atomic mass unit, 10^{-3} kg mol ⁻¹ N_A^{-1}	u	1.660 540 2	$\times 10^{-27}$ kg	0.17
Faraday constant	$F (= N_A e)$	9.648 533 83	$\times 10^4$ C mol ⁻¹	0.086
Newtonian constant of gravitation	G	6.6742	$\times 10^{-11}$ N m ² Kg ⁻²	150

Spectroscopy and atoms

Planck constant	h	4.135 667 43	$\times 10^{-15}$ eV s	0.085
Planck constant ($h/2\pi$)	\hbar	6.582 119 15	$\times 10^{-16}$ eV s	0.085
Charge/mass ratio of electron	$-e/m_e$	-1.758 820 12	$\times 10^{11}$ C kg ⁻¹	0.086
Fine structure constant	α	7.297 352 568	$\times 10^{-3}$	0.0033
Fine structure constant, reciprocal	α^{-1}	137.035 999 11		0.0033
Rydberg constant (fixed nucleus)	R_∞	10 973 731.568 525	m ⁻¹	0.0000066
Bohr radius ($4\pi/\mu_0 c^2$) $\hbar^2/m_e e^2$	a_0	5.291 772 108	$\times 10^{-11}$ m	0.00333
Compton wavelength of electron	λ_c	2.426 310 238	$\times 10^{-12}$ m	0.0067
Compton wavelength of electron $\div 2\pi$	$\hbar/m_e c$	3.861 592 678	$\times 10^{-13}$ m	0.0067
Classical 'radius' of electron ($\mu_0 c^2/4\pi$) e^2/mc^2	r_e	2.817 940 325	$\times 10^{-15}$ m	0.01
Thomson cross-section $8\pi r_e^2/3$	σ_e	6.652 458 73	$\times 10^{-29}$ m ²	0.02
Zeeman effect, μ_B/hc		46.686 4507	m ⁻¹ T ⁻¹	0.086
Bohr magneton $e\hbar/2m_e$	μ_B	9.274 015 4	$\times 10^{-24}$ J T ⁻¹	0.086
Nuclear magneton $e\hbar/2m_p$	μ_N	5.050 783 43	$\times 10^{-27}$ J T ⁻¹	0.086
Ratio of masses proton/electron	m_p/m_e	1 836.152 672 61		0.00046
Gyromagnetic ratio of proton	$\gamma_p = \mu_p / \frac{1}{2}\hbar$	2.675 222 05	$\times 10^8$ s ⁻¹ T ⁻¹	0.086
in H ₂ O, sph., 25 °C	γ'_p	2.675 153 33	$\times 10^8$ s ⁻¹ T ⁻¹	0.086
in H ₂ O (cycles), sph., 25 °C	$\gamma'_p/2\pi$	4.257 638 75	$\times 10^7$ Hz T ⁻¹	0.086

Conversion factors for mass, energy and wavelength

Energy and mass

Electron volt		1.602 176 53	$\times 10^{-19}$ J	0.17
Atomic mass unit		931.494 043	MeV	0.086

1 MeV		1.073 544 171	$\times 10^{-3}$	u	0.0086
Rest-mass of electron		0.510 998 9186		MeV	0.17
1 eV per molecule		9.648 533 83	$\times 10^7$	J kmol ⁻¹	0.086

Frequency, wavelength and energy

Quantum energy ÷ wave number		1.986 445 61	$\times 10^{-25}$	J m	0.18
Energy × wavelength		1.239 841 91	$\times 10^{-6}$	eV m	0.0085
Wave number ÷ energy		8.065 544 45	$\times 10^5$	eV ⁻¹ m ⁻¹	0.085
Quantum energy ÷ frequency .		4.135 667 43	$\times 10^{-15}$	eV Hz ⁻¹	0.0085
Frequency ÷ energy		2.417 989 40	$\times 10^{14}$	Hz eV ⁻¹	0.085

Thermal constants

Molar gas constant	R	8.314 472		J mol ⁻¹ K ⁻¹	1.7
Loschmidt constant (number of molecules in 1 m ³ of ideal gas at stp)	n_0	2.686 7773	$\times 10^{25}$	m ⁻³	1.8
Boltzmann constant R/N_A	k	1.380 6505	$\times 10^{-23}$	J K ⁻¹	1.8
Boltzmann constant		8.617 343	$\times 10^{-5}$	eV K ⁻¹	1.8
Stefan-Boltzmann constant (σ) [†]	$2\pi^5 k^4/15c^2 h^3$	5.670 400	$\times 10^{-8}$	Wm ⁻² K ⁻⁴	7
Constant in Planck formula (c_1) [†]	$2\pi h c^2$	3.741 771 38	$\times 10^{-16}$	W m ²	0.17
Constant in Planck formula (c_2) [†]	hc/k	1.438 7752	$\times 10^{-2}$	m K	1.7

[†] see also section 2.5.2.

Note: Magnetic moments are defined so that mechanical energy = $-\mu \cdot \mathbf{B}$; the unit is $1 \text{ J T}^{-1} = 1 \text{ J Wb}^{-1} \text{ m}^2 = 1 \text{ A m}^2$ (1 tesla = 1 weber m⁻² \equiv 10⁴ gauss).

*Standard values**

Von Klitzing constant R_{K-90} 25 812.807 ($1 \pm 0.2 \times 10^{-6}$) Ω
Josephson constant K_{J-90} 483 579.9 ($1 \pm 0.4 \times 10^{-6}$) GHz/V

* These standard values were recommended by the CCE in 1989 for adoption from 1.1.1990. The CCE considered later data than were available to Cohen and Taylor (1987) and they are not necessarily identical with h/e^2 and $2e/h$ respectively. They were defined by the CCE in order to help ensure global uniformity of standards of resistance and emf. (The suffix -90 is used here to indicate that they may be revised at a later date, although it is usually omitted.)

Reference

P. J Mohr and B. N. Taylor, Rev. Mod Phys., **76**, no 4(Oct 2004)

B.W.Petley

Reviewed: 11 Jan 2005

[« Previous Subsection](#)

[Next Subsection »](#)

[Home](#) | [About](#) | [Table of Contents](#) | [Advanced Search](#) | [Copyright](#) | [Feedback](#) | [Privacy](#) | [^ Top of Page ^](#)

This site is hosted and maintained by the [National Physical Laboratory](#) © 2011.